
Specimen Collection & Handling


Follow Order of Draw and process all specimens within 2 hours of collection.

Order of Draw	Type/Additive	Invert Immediately	Centrifuge	Refrigerate	Common Tests
AND	Bactec [™] Culture Bottles	×	×	×	Blood Cultures, Fluid Cultures
Son Burgardon and Burgardon and Company an					
Orne CC Vacas CS (disposance or grant of the control of the contro	Royal Blue Trace Element K₂EDTA	8-10 times	×	✓	Aluminum, Zinc
De de la constante de la const	Red/Gray Stopper No additive	×	×	×	Discard tube for Light Blue tube, DO NOT send.*
	Light Blue Sodium Citrate	4 times	×	×	Prothrombin Time (PT)
	Red Clot activator	5-10 times	After clotting	✓	Therapeutic Drugs, Carnitine
William Annual Control of the Contro	Green Lithium Heparin	8-10 times	Immediately	✓	Routine Chemistry, Hepatitis, PTH, Ferritin, HIV, SARS-CoV-2 Ab, Vitamin D
Francisco Franci	Lavender K₂EDTA	8-10 times	X	✓	Hematology, A1C, Fluid Cell Count, some Reflex testing


^{*}Only use if Light Blue tube is first in order of draw.



Labeling & Handling











Room