

## CALCULATIONS

<b>Adjusted Calcium (mg/dL)</b>	Total Calcium + 0.8 x (4.0-Albumin)
<i>Included In</i>	Adjusted Calcium Phosphorus Product Comprehensive Metabolic Panel Comprehensive Metabolic Panel w/Phosphorus
<i>Note</i>	Calculation provided only when Albumin is <4.0 g/dL

<b>Adjusted Calcium Phosphorus Product (mg<sup>2</sup>/dL<sup>2</sup>)</b>	(Total Calcium (mg/dL) + 0.8 x (4.0-Albumin (g/dL))) x Phosphorus
<i>Included In</i>	Adjusted Calcium Phosphorus Product Comprehensive Metabolic Panel w/Phosphorus
<i>Note</i>	Calculation provided only when Albumin is <4.0 g/dL

<b>A/G Ratio</b>	Albumin/Globulin
<i>Included In</i>	Comprehensive Metabolic Panel Comprehensive Metabolic Panel w/Phosphorus

<b>Anion Gap (mEq/L)</b>	Sodium – (Chloride + CO <sub>2</sub> )
<i>Included In</i>	Basic Metabolic Panel Comprehensive Metabolic Panel Comprehensive Metabolic Panel w/ Phosphorus Electrolytes Renal Function Panel

<b>Calcium Phosphorus Product (mg<sup>2</sup>/dL<sup>2</sup>)</b>	Total Calcium x Phosphorus
<i>Included In</i>	Calcium Phosphorus Product Adjusted Calcium Phosphorus Product Comprehensive Metabolic Panel w/Phosphorus

<b>estimated Glomerular Filtration Rate (eGFR) Creatinine (mL/min/1.73 m<sup>2</sup>)</b>	$142 \times \min(\text{Scr}/K, 1)^\alpha \times \max(\text{Scr}/K, 1)^{-1.200} \times 0.9938^{\text{Age}} \times 1.012$ [if female]
<i>Note</i>	Estimated GFR (eGFR) using CKD-EPI 2021 where Scr = standardized creatinine in mg/dL K = 0.7 (females) or 0.9 (males) $\alpha = -0.241$ (females) or $-0.302$ (males) min = indicates the minimum of Scr/K or 1 max = indicates the maximum of Scr/K or 1 Age = years

<b>estimated Glomerular Filtration Rate (eGFR) Creatinine-Cystatin C (mL/min/1.73 m<sup>2</sup>)</b>	$135 \times \min(\text{Scr}/K, 1)^\alpha \times \max(\text{Scr}/K, 1)^{-0.544} \times \min(\text{Scys}/0.8, 1)^{-0.323} \times \max(\text{Scys}/0.8, 1)^{-0.778} \times 0.9961^{\text{Age}} \times 0.963$ [if female]
<i>Note</i>	Estimated GFR (eGFR) using CKD-EPI 2021 where Scr = standardized creatinine in mg/dL K = 0.7 (females) or 0.9 (males) $\alpha = -0.241$ (females) or $-0.302$ (males) min = indicates the minimum of Scr/K or 1 max = indicates the maximum of Scr/K or 1 Scys = standardized Cystatin C in mg/L Age = years

<b>Globulin (g/dL)</b>	Total Protein–Albumin
<i>Included In</i>	Comprehensive Metabolic Panel Comprehensive Metabolic Panel w/Phosphorus

## CALCULATIONS

<b>Hemoglobin x 3 (g/dL)</b>	Hemoglobin x 3
<i>Included In</i>	Complete Blood Count (CBC) & Differential Complete Blood Count (CBC) & Differential w/Reticulocytes Hemoglobin Hemoglobin & Hematocrit (H&H) Hemogram (Complete Blood Count w/o Differential)

<b>Iron Status w/Iron &amp; Transferrin</b>	
Total Iron Binding Capacity (TIBC) (µg/dL)	Transferrin x 1.4
Transferrin Saturation (%)	(Iron/(Transferrin x 1.4)) x 100

<b>Lipid Panel (Coronary Risk Profile)</b>	
Cholesterol/HDL Ratio	Cholesterol/HDL
Low Density Lipoprotein (mg/dL)	Cholesterol – (Very Low Density Lipoprotein + HDL)
Very Low Density Lipoprotein (mg/dL)*	Triglycerides/5
Note*	Only provided if Triglyceride is <400 mg/dL

<b>% Recirculation</b>	(Systemic BUN – Arterial BUN)/(Systemic BUN – Venous BUN) x 100
<i>Included In</i>	Recirculation Study

<b>Prothrombin Time (Protime)</b>	
INR	(PT Ratio) <sup>ISI</sup> PT Ratio = (Patient PT/Mean Normal PT) <sup>ISI</sup> Mean Normal PT = Geometric Mean ISI = International Sensitivity Index

## CALCULATIONS

HEMODIALYSIS CALCULATIONS	
<b>Kt/V Equilibrated (eqKt/V)</b>	$(0.924 \times \ln \text{Kt/V}) - ((0.395 \times \ln \text{Kt/V}) / (\text{Min} / 60)) + 0.056$
Included In	Kt/V Standard, URR Kt/V Standard, Natural Log, URR
Note	Leypoldt Formula For patient dialyzing 2 or 4-6 times per week For Kt/V Standard calculation purposes only, not reported
<b>Kt/V Jindal</b>	$(0.04 \times ((\text{Pre BUN} - \text{Post BUN}) / \text{Pre BUN} \times 100) - 1.2)$
Note	Jindal Formula (Not KDOQI recommended) The HD Adequacy Work Group feels this formula should not be used to measure delivered dose of Hemodialysis. (K/DOQI Clinical Practice Guidelines for Hemodialysis Adequacy: Update 2000, Guideline 2)
<b>Kt/V Natural Log (lnKt/V)</b>	$(-\ln((\text{Post BUN}/\text{Pre BUN}) - (0.008 \times \text{Treatment Time in mins}/60)) + ((4 - (3.5 \times (\text{Post BUN}/\text{Pre BUN})) \times (\text{Pre WT} - \text{Post WT})/\text{Post WT}))$
Included In	Kt/V Natural Log, URR Kt/V Natural Log, URR, nPNA Kt/V Standard, Natural Log, URR
Note	Daugirdas II Formula The K/DOQI recommendations are: Prescribed dose of hemodialysis: Kt/V of 1.3 Delivered dose of hemodialysis: Kt/V >1.2
<b>Kt/V Standard (stdKt/V)</b>	$(168 \times (1 - \exp(-\text{eqKt/V})) / (\text{Min}/60)) / ((1 - \exp(-\text{eqKt/V})) / \text{eqKt/V} + (168/\text{Number of Treatment}/(\text{Min}/60)) - 1)$
Included In	Kt/V Standard, URR Kt/V Standard, Natural Log, URR
Note	Leypoldt Formula For patient dialyzing 2 or 4-6 times per week
<b>Kt/V Residual</b>	$(\text{Urine Urea Nitrogen}/\text{Blood BUN}) \times (\text{Urine Volume}/\text{Urine Collection Time}) \times (10.08/\text{VSA})$
Included In	Kt/V Natural Log, URR Kt/V Natural Log, URR, nPNA Kt/V Standard, URR Kt/V Standard, Natural Log, URR
Note	Only calculated if urine provided; added to Kt/V Natural Log or Standard up to 90 days
<b>nPNA, Hemodialysis</b>	<ol style="list-style-type: none"> <li>Treatment #1: Beginning of week PNA (PCR) = <math>\text{Pre BUN} / (36.3 + 5.48 \times \text{Kt/V Natural Log} + 53.5 / \text{Kt/V Natural Log}) + 0.168</math></li> <li>Treatment #2: Midweek PNA (PCR) = <math>\text{Pre BUN} / (25.8 + 1.15 \times \text{Kt/V Natural Log} + 56.4 / \text{Kt/V Natural Log}) + 0.168</math></li> <li>Treatment #3: End of week PNA (PCR) = <math>\text{Pre BUN} / (16.3 + 4.3 \times \text{Kt/V Natural Log} + 56.6 / \text{Kt/V Natural Log}) + 0.168</math></li> </ol>
Included In	Kt/V Natural Log, URR, nPNA
Note	nPNA calculation is only applicable to patients on thrice-weekly dialysis without significant residual function. nPNA calculated from Kt/V without formal kinetic modeling according to Depner T and Daugirdas J: JASN 1996;7:780-785.
<b>Urea Reduction Ratio (%)</b>	$(1 - (\text{Post BUN}/\text{Pre BUN})) \times 100$
Included In	Kt/V Jindal (Not K/DOQI Recommended) Kt/V Natural Log, URR Kt/V Natural Log, URR, nPNA Kt/V Standard, Natural Log, URR Kt/V Standard, URR Urea Reduction Ratio w/Pre and Post BUN
<b>Ultrafiltration Rate (UFR) (mL/kg/hr)</b>	$((\text{pre-weight} - \text{post-weight}) \times 1000) / (\text{delivered time in mins}/60) / \text{post-weight in kg}$
Included In	Kt/V Natural Log, URR Kt/V Natural Log, URR, nPNA Kt/V Standard, Natural Log, URR Kt/V Standard, URR

## CALCULATIONS

PD ADEQUACY CALCULATIONS	
<b>Weekly Total Kt/V</b>	Weekly Residual Kt/V + Weekly Dialysate Kt/V
<b>Weekly Residual Kt/V</b>	$((\text{Urine Urea Nitrogen/BUN}) \times (\text{Urine Volume (mL)/Urine Collection Time (min)}) \times 10.08) / \text{VSA}$
<i>Note</i>	<i>Calculated if urine sample provided</i>
<b>Weekly Dialysate Kt/V</b>	$((\text{Dialysate Urea Nitrogen/BUN}) \times (24 \text{ Hour Dialysate Drain Volume (mL)/1000}) \times 7) / \text{VSA}$
<b>Weekly Total CrCl (L/wk/1.73 m<sup>2</sup>)</b>	Weekly Residual GFR + Weekly Dialysate Creatinine Clearance
<b>Weekly Residual GFR (L/wk/1.73 m<sup>2</sup>)</b>	Arithmetic Mean of Weekly Urea Clearance and Weekly Creatinine Clearance $((\text{Urine Urea Nitrogen/BUN}) \times (\text{Urine Volume (mL)/Urine Collection Time (min)}) \times 10.08) + (\text{Urine Creatinine/Plasma Creatinine} \times \text{Urine Volume (mL)/Urine Collection Time (min)} \times 10.08) / 2 \times 1.73 / \text{BSA}$
<b>Weekly Dialysate CrCl (L/wk/1.73 m<sup>2</sup>)</b>	$(\text{Dialysate Corrected Creatinine/Plasma Creatinine}) \times (24 \text{ Hour Dialysate Drain Volume (mL)/1000}) \times 7 \times 1.73 / \text{BSA}$
<b>Weekly Residual CrCl (L/wk/1.73 m<sup>2</sup>)</b>	$(\text{Urine Creatinine/Blood Creatinine}) \times (\text{Urine Volume (mL)/Urine Collection Time (min)}) \times (1.73 / \text{BSA}) \times 10.08$
<b>Creatinine Clearance (mL/min/1.73m<sup>2</sup>)</b>	$(\text{Urine Creatinine/Blood Creatinine}) \times (\text{Urine Volume (mL)/Urine Collection Time (min)}) \times (1.73 / \text{BSA})$
<b>Corrected Creatinine, 24 Hour (mg/dL)</b>	Creatinine at 24 Hour Dwell – (Glucose at 24 Hour Dwell x 0.00010386)
<b>nPNA, Peritoneal Dialysis (g/kg/day)</b>	$(10.76 \times ((0.69 \times \text{UNA}) + 1.46)) / (\text{VSA}/0.58)$
<b>Protein Nitrogen Appearance (PNA) (g/day)</b>	$10.76 \times ((0.69 \times \text{UNA}) + 1.46)$
<b>UNA (g/day)</b>	$(24 \text{ Hour Drain Volume (mL)} \times 24 \text{ Hour Urea Dialysate}) / 100000 + (\text{Urine Volume (mL)} \times \text{Urine Urea Nitrogen}) / 100000 \times (1440 / \text{Total Urine Collection Time (min)})$
<i>Note</i>	<i>UNA used for PNA calculation purposes only, not reported</i>
<b>Body Surface Area (BSA) (m<sup>2</sup>)</b>	Adult (≥16 years) uses DuBois and DuBois formula $\text{BSA (m}^2) = 0.007184 \times \text{Wt}^{0.425} \times \text{Ht}^{0.725}$ Pediatric (< 16 years) uses Haycock formula $\text{BSA (m}^2) = 0.024265 \times \text{Wt}^{0.5378} \times \text{Ht}^{0.3964}$ where weight (Wt) is in kilograms and height (Ht) is in centimeters
<b>Volume from Surface Area (VSA) (Liters)</b>	Adult (≥16 years) uses Hume and Weyers formula Male: $V = -14.012934 + 0.296785 \times \text{Wt} + 0.194786 \times \text{Ht}$ Female: $V = -35.270121 + 0.183809 \times \text{Wt} + 0.344547 \times \text{Ht}$ Pediatric (<16 years) uses Friis-Hansen formula $V = 0.135 \times \text{Wt}^{0.666} \times \text{Ht}^{0.535}$ where weight (Wt) is in kilograms and height (Ht) is in centimeters

## CALCULATIONS

<b>FLUID CALCULATIONS</b>	
<b>Peritoneal Equilibration Test (PET) Fast</b>	
Corrected Creatinine, 4 Hour (mg/dL)	Creatinine at 4 Hour Dwell – (Glucose at 4 Hour Dwell x 0.00010386)
Corrected Creatinine D/P, 4 Hour	Corrected Creatinine at 4 Hour Dwell/Plasma Creatinine
<b>Peritoneal Equilibration Test (PET) Standard</b>	
Corrected Creatinine, 0 Hour, 2 Hour, 4 Hour (mg/dL)	Creatinine at 0 or 2 or 4 Hour Dwell – (Glucose at 0 or 2 or 4 Hour Dwell x 0.00010386)
Corrected Creatinine D/P, 0 Hour	Corrected Creatinine at 0 Hour Dwell/Plasma Creatinine
Corrected Creatinine D/P, 2 Hour	Corrected Creatinine at 2 Hour Dwell/Plasma Creatinine
Corrected Creatinine D/P, 4 Hour	Corrected Creatinine at 4 Hour Dwell/Plasma Creatinine
Glucose D/D0, 2 Hour	Glucose at 2 Hour Dwell/Glucose at 0 Hour Dwell
Glucose D/D0, 4 Hour	Glucose at 4 Hour Dwell/Glucose at 0 Hour Dwell
Urea D/P, 0 Hour	Urea at 0 Hour Dwell/Plasma Urea
Urea D/P, 2 Hour	Urea at 2 Hour Dwell/Plasma Urea
Urea D/P, 4 Hour	Urea at 4 Hour Dwell/Plasma Urea
<b>Peritoneal Equilibration Test (PET) Modified</b>	
Corrected Creatinine, 0 Hour, 1 Hour, 2 Hour, 4 Hour (mg/dL)	Creatinine at 0 or 1 or 2 or 4 Hour Dwell – (Glucose at 0 or 1 or 2 or 4 Hour Dwell x 0.00010386)
Corrected Creatinine D/P, 4 Hour	Corrected Creatinine at 4 Hour Dwell/Plasma Creatinine
Sodium D/P, 0 Hour	Sodium at 0 Hour Dwell / Plasma Sodium
Sodium D/P, 1 Hour	Sodium at 1 Hour Dwell / Plasma Sodium
Sodium D/P, 2 Hour	Sodium at 2 Hour Dwell / Plasma Sodium
Sodium D/P, 4 Hour	Sodium at 4 Hour Dwell / Plasma Sodium
<b>Fluid, 24-Hour Dwell</b>	
Corrected Creatinine, 24 Hour (mg/dL)	Creatinine at 24 Hour Dwell – (Glucose at 24 Hour Dwell x 0.00010386)
<b>Fluid, Overnight Dwell</b>	
Corrected Creatinine, Overnight (mg/dL)	Creatinine Overnight Dwell – (Glucose Overnight Dwell x 0.00010386)
<b>Gotch PD QA</b>	
Corrected Creatinine, PD QA (mg/dL)	(Corrected Creatinine, PD QA – Glucose PD QA x 0.00010386)
Total Protein, PD QA (g/dL)	Total Protein, PD QA / 1000
<b>Gotch PD Exchange 1</b>	
Corrected Creatinine, PD Exchange 1 (mg/dL)	(Corrected Creatinine, PD Exchange 1 – Glucose PD Exchange 1 x 0.00010386)
Total Protein, PD Exchange 1 (g/dL)	Total Protein, PD Exchange 1 / 1000
<b>Gotch PD Exchange 2</b>	
Corrected Creatinine, PD Exchange 2 (mg/dL)	(Corrected Creatinine, PD Exchange 2 – Glucose PD Exchange 2 x 0.00010386)
Total Protein, PD Exchange 2 (g/dL)	Total Protein, PD Exchange 2 / 1000
<b>Gotch PD Exchange 3</b>	
Corrected Creatinine, PD Exchange 3 (mg/dL)	(Corrected Creatinine, PD Exchange 3 – Glucose PD Exchange 3 x 0.00010386)
Total Protein, PD Exchange 3 (g/dL)	Total Protein, PD Exchange 3 / 1000
<b>Gotch PD Exchange 4</b>	
Corrected Creatinine, PD Exchange 4 (mg/dL)	(Corrected Creatinine, PD Exchange 4 – Glucose PD Exchange 4 x 0.00010386)
Total Protein, PD Exchange 4 (g/dL)	Total Protein, PD Exchange 4 / 1000
<b>Gotch PD Exchange 5</b>	
Corrected Creatinine, PD Exchange 5 (mg/dL)	(Corrected Creatinine, PD Exchange 5 – Glucose PD Exchange 5 x 0.00010386)
Total Protein, PD Exchange 5 (g/dL)	Total Protein, PD Exchange 5 / 1000

## CALCULATIONS

<b>URINE CALCULATIONS</b>	
<b>24 Hour Urine Creatinine Clearance (Residual Renal Creatinine Clearance)</b>	
Creatinine Clearance (mL/min/1.73m <sup>2</sup> )	$(\text{Urine Creatinine}/\text{Blood Creatinine}) \times (\text{Urine Volume (mL)}/\text{Urine Collection Time (min)}) \times (1.73/\text{BSA})$
Body Surface Area (BSA)	Adult ( $\geq 16$ years) uses DuBois and DuBois formula $\text{BSA (m}^2\text{)} = 0.007184 \times \text{Wt}^{0.425} \times \text{Ht}^{0.725}$ Pediatric (< 16 years) uses Haycock formula $\text{BSA (m}^2\text{)} = 0.024265 \times \text{Wt}^{0.5378} \times \text{Ht}^{0.3964}$ where weight (Wt) is in kilograms and height (Ht) is in centimeters

<b>Residual Urea Clearance, KrU – for Hemodialysis only</b>	
KrU (mL/min)	$(\text{Urine Urea Nitrogen} \times \text{Urine Volume (mL)}) / (\text{Blood BUN} \times 0.9 \times \text{Total Urine Collection Time (min)})$
Kt/V Residual	$(\text{Urine Urea Nitrogen}/\text{Blood BUN}) \times (\text{Urine Volume (mL)}/\text{Urine Collection Time (min)}) \times (10.08/\text{VSA})$
Volume from Surface Area (VSA) (Liters)	Adult ( $\geq 16$ years) uses Hume and Weyers formula Male: $V = -14.012934 + 0.296785 \times \text{Wt} + 0.194786 \times \text{Ht}$ Female: $V = -35.270121 + 0.183809 \times \text{Wt} + 0.344547 \times \text{Ht}$ Pediatric (<16 years) uses Friis-Hansen formula $V = 0.135 \times \text{Wt}^{0.666} \times \text{Ht}^{0.535}$ where weight (Wt) is in kilograms and height (Ht) is in centimeters

<b>24 Hour Urine Creatinine</b>	
Urine Creatinine, 24 Hour (mg/24 hr)	$((\text{Urine Creatinine in mg/dL} \times \text{Urine Volume in mL})/100) \times (1440 / \text{Total Urine Collection Time in min})$

<b>24 Hour Urine Urea Nitrogen</b>	
Urine Urea Nitrogen, 24 Hour (g/24 hr)	$((\text{Urine Urea Nitrogen in mg/dL}/100) \times (\text{Urine Volume in mL}/(\text{Total Urine Collection Time in mins}/1440)))/1000$

<b>24 Hour Urine Total Protein with Creatinine</b>	
Urine Protein, 24 Hour (mg/24 hr)	$(\text{Urine Total Protein}/100) \times ((\text{Urine Volume in mL})/(\text{Total Urine Collection Time in mins}/1440))$
Urine Total Protein/ Creatinine Ratio, 24 Hour (mg/g creat)	$((\text{Urine Total Protein}/100) \times ((\text{Urine Volume in mL})/(\text{Total Urine Collection Time in mins}/1440)))/(((\text{Urine Creatinine}/100) \times (\text{Urine Volume in mL}/(\text{Total Urine Collection Time in mins}/1440))))/1000$
Urine Creatinine, 24 Hour (g/24 hr)	$((\text{Urine Creatinine}/100) \times (\text{Urine Volume in mL}/(\text{Total Urine Collection Time in mins}/1440)))/1000$

<b>Random Urine Total Protein with Creatinine</b>	
Urine Total Protein/ Creatinine Ratio (mg/g creat)	$(\text{Total Protein, Random Urine}/\text{Creatinine, Random Urine}) \times 1000$